

JACK POLYNOMIALS AND HILBERT SCHEMES OF POINTS ON SURFACES

HIRAKU NAKAJIMA

1. INTRODUCTION

The Jack (symmetric) polynomials $P_\lambda^{(\alpha)}(x)$ form a class of symmetric polynomials which are indexed by a partition λ and depend rationally on a parameter α . They reduced to the Schur polynomials when $\alpha = 1$, and to other classical families of symmetric polynomials for several specific parameters. Recently they attracts attention from various points of view, for example the integrable systems and combinatorics. They are simultaneous eigenfunctions of certain commuting families of differential operators, appearing in the integrable system called the Calogero-Sutherland system (see e.g., [2] and the reference therein). On the other hand, Macdonald studied their combinatorial properties. In fact, he introduced an even more general class of symmetric functions (a two parameter family), which have many common combinatorial features as Jack polynomials (see [17, Chapter 6]).

It is well-known that Schur polynomials can be realized as certain elements of homology groups of Grassmann manifolds (see e.g., [9, Chapter 14]). The purpose of this paper is to give a similar geometric realization for Jack polynomials. However, spaces which we use are totally different. Our spaces are Hilbert schemes of points on a surface X which is the total space of a line bundle L over the projective line $\mathbb{CP}^1$. The parameter α in Jack polynomials relates to our surface X by

$$(1.1) \quad \alpha = -\langle C, C \rangle = -c_1(L)[C],$$

where C is the zero section, and $\langle C, C \rangle$ is the self-intersection number of C . It seems difficult to realize Jack polynomials in homology groups of Grassmann manifolds since they have no parameter.

The Hilbert scheme $X^{[n]}$ parametrizing 0-dimensional subschemes of length n on X has been studied by various peoples (see [11] and the reference therein). The author and Grojnowski independently showed that the direct sum of the homology groups of $X^{[n]}$ (the summation is over n) is a representation space of the Heisenberg algebra (boson Fock space) [13, 18]. It is also well-known that the space of symmetric polynomials is the same representation space. This is the relationship between symmetric functions and Hilbert schemes.

1991 *Mathematics Subject Classification*. Primary 14C05, 05E05.

Key words and phrases. Jack polynomials, Hilbert scheme.

Author supported in part by Grant-in-Aid for Scientific Research (No. 08740010), Ministry of Education, Science and Culture, Japan.

Now we explain our realization in more detail. There are two characterizations of Jack polynomials:

- (a) an orthogonal basis such that the transition matrix to monomial symmetric functions is strictly upper triangular (see Theorem 2.3), or
- (b) simultaneous eigenfunctions of a family of commuting differential operators (see above).

We use the characterization (a). We first identify the complexified ring of symmetric functions with the direct sum of the middle degree homology groups of Hilbert schemes as above. We then identify the inner product with the intersection pairing (Theorem 3.5). This result was essentially proved in [13, 18] combined with [7]. Then monomial symmetric functions are identified with fundamental classes of certain middle dimensional subvarieties in the Hilbert schemes (Theorem 4.6), which were first introduced by Grojnowski [13]. Finally, in order to get mutually orthogonal elements, we use the *localization*. We define an S^1 -action on the surface X which induces an action on the Hilbert scheme. The localization, which goes back to Bott residue formula [3], enables computations of the intersection product to be reduced to the fixed point set of the S^1 -action. Then cohomology classes which localize to different fixed point components are mutually orthogonal. This is the mechanism to construct an orthogonal basis. (In this paper, we shall use the equivariant cohomology to formulate the localization following [1].) The relation to the subvarieties corresponding to monomial symmetric functions can be studied as follows. For each fixed point component, we associate a locally closed submanifold (“stratum”) consisting of points which converge to the fixed point components when they moved by the $\mathbb{C}^*$ -actions which extends the S^1 -action (see (4.4)). The dominance order is identified with the closure relation of the strata (4.13). The subvarieties corresponding monomial symmetric functions are the closures of stratum. Then it is easy to check that the transition matrix is strictly upper triangular.

In fact, almost all arguments work for the total space of a line bundle over any compact Riemann surface. We use the middle degree homology group of Hilbert schemes on which only $H^2(C)$ contributes. ($H^0(C)$ and $H^1(C)$ contribute only to lower degrees.)

The motivation of this work comes from Wilson’s observation [20] that “completed phase space” for the complex Calogero-Moser system (a cousin of the Calogero-Sutherland system) is diffeomorphic to the Hilbert scheme of points on the affine plane. (The author learned this observation from Segal’s talk at Warwick, 1996 March.) This made the author to look for the connection between Hilbert schemes and Jack polynomials. However, the connection found in this paper does not follow the Wilson’s route. Clarifying relation between these two connections should be an important problem. For example, it is desirable to have a geometric realization of commuting differential operators.

It seems natural to conjecture that we get similar geometric realization of Macdonald polynomials (two parameter family explained above) if we replace the homology by the equivariant K-theory. Analogous phenomenon was found for the affine Weyl group and the affine Hecke algebra. The group ring of the affine Weyl group is realized on the homology

group of the cotangent bundle of the flag manifold, while the Hecke algebra is realized on the equivariant K-theory. (See [5].) We hope to return back in near future.

Acknowledgment. The author would like to thank K. Hasegawa who told him the relation between Jack polynomials and the Calogero-Sutherland system just after Segal's talk at Warwick. It is also a pleasure to acknowledge discussions with T. Gocho, A. Matsuo and H. Ochiai during the seminar on Hilbert schemes of points held in 1996 spring. In particular, Matsuo's talk on Macdonald polynomials was very helpful.

2. PRELIMINARIES

In this section we review the theory of symmetric functions and the equivariant cohomology for later use.

2(i). Symmetric Functions. First we briefly recall the theory of symmetric functions. See [17] for detail.

A *partition* $\lambda = (\lambda_1, \lambda_2, \lambda_3, \dots)$ is a nonincreasing sequence of nonnegative integers such that $\lambda_i = 0$ for all but finitely many i . Let $|\lambda| = \sum_i \lambda_i$. We say λ is a partition of n if $|\lambda| = n$. We also use another presentation $\lambda = (1^{m_1} 2^{m_2} \dots)$ where $m_k = \#\{i \mid \lambda_i = k\}$. Number of nonzero entries in λ is called *length* of λ and denoted by $l(\lambda)$.

If λ and μ are partitions, we define $\lambda \cup \mu$ be the partition whose entries are those of λ and μ arranged in the descending order.

For a partition $\lambda = (\lambda_1, \lambda_2, \dots)$, we give a Young diagram such that the number of boxes in the i th column is λ_i . Remark that our convention differs from one used in [17]. Our diagram is rotated by $\pi/2$ from one used in [17].

The conjugate of a partition λ is the partition λ' whose diagram is the transpose of the diagram of λ , i.e.,

$$(2.1) \quad \lambda'_i = \#\{j \mid \lambda_j \geq i\}.$$

We define $\lambda \geq \mu$ if $|\lambda| = |\mu|$ and

$$(2.2) \quad \lambda_1 + \lambda_2 + \dots + \lambda_i \geq \mu_1 + \mu_2 + \dots + \mu_i \quad \text{for all } i.$$

This defines a partial order on the set of partitions and is called *dominance order*. Note that $\lambda \geq \mu$ if and only if $\mu' \geq \lambda'$.

Let Λ_N be the ring of symmetric functions

$$\Lambda_N = \mathbb{Z}[x_1, \dots, x_N]^{\mathfrak{S}_N},$$

where the symmetric group $\mathfrak{S}_N$ acts by the permutation of the variables. It is a graded ring:

$$\Lambda_N = \bigoplus_{n \geq 0} \Lambda_N^n,$$

where Λ_N^n consists of the homogeneous symmetric functions of degree n . It is more relevant for us to consider symmetric functions in “infinitely many variables” formulated as follows:

let $M \geq N$ and consider the homomorphism

$$\mathbb{Z}[x_1, \dots, x_M] \rightarrow \mathbb{Z}[x_1, \dots, x_N]$$

which sends $x_{N+1}, \dots, x_M$ to 0. We have induced homomorphisms

$$\rho_{M,N}^n: \Lambda_M^n \rightarrow \Lambda_N^n,$$

which is surjective for any $M \geq N$, and bijective for $M \geq N \geq n$. Let

$$\Lambda^n \stackrel{\text{def.}}{=} \varprojlim \Lambda_N^n.$$

Then the ring of symmetric functions in infinitely many variables is defined by

$$\Lambda \stackrel{\text{def.}}{=} \bigoplus_n \Lambda^n.$$

In the relationship between symmetric functions and Hilbert schemes, the degree n corresponds the number of points, while the number of variables N are irrelevant. This is the only reason why we use the different notation from [17].

There are several distinguished classes of symmetric functions. The first class is the *monomial symmetric function* m_λ . Let λ be a partition with $l(\lambda) \leq N$. Let

$$m_\lambda(x_1, \dots, x_N) \stackrel{\text{def.}}{=} \sum_{\alpha \in \mathfrak{S}_N \cdot \lambda} x_1^{\alpha_1} \cdots x_N^{\alpha_N} = \frac{1}{\#\{\sigma \in \mathfrak{S}_N \mid \sigma \cdot \lambda = \lambda\}} \sum_{\sigma \in \mathfrak{S}_N} x_1^{\lambda_{\sigma(1)}} \cdots x_N^{\lambda_{\sigma(N)}},$$

where $\alpha = (\alpha_1, \dots, \alpha_N)$ runs over all distinct permutation of $(\lambda_1, \lambda_2, \dots, \lambda_N)$. If $l(\lambda) \leq N$, we have

$$\rho_{M,N}^n m_\lambda(x_1, \dots, x_M) = m_\lambda(x_1, \dots, x_N).$$

Hence m_λ defines an element in Λ , which is also denoted by m_λ . Then $\{m_\lambda\}_\lambda$ is a basis for Λ .

The n th power sum is

$$p_n \stackrel{\text{def.}}{=} \sum x_i^n = m_{(n)}.$$

For a partition $\lambda = (\lambda_1, \lambda_2, \dots)$, let $p_\lambda = p_{\lambda_1} p_{\lambda_2} \cdots$. Then $\{p_\lambda\}_\lambda$ is a basis for $\Lambda \otimes \mathbb{Q}$. (It is *not* a $\mathbb{Z}$ -basis for Λ .)

For a positive real number α , we define an inner product $\langle \cdot, \cdot \rangle$ on $\Lambda \otimes \mathbb{Q}$ by

$$\langle p_\lambda, p_\mu \rangle \stackrel{\text{def.}}{=} \alpha^{l(\lambda)} z_\lambda \delta_{\lambda\mu},$$

where $z_\lambda = \prod k^{m_k} m_k!$ for $\lambda = (1^{m_1} 2^{m_2} \cdots)$.

The Jack polynomials are defined by

Theorem 2.3 ([17, 10.13]). *For each partition λ , there is a unique symmetric polynomial $P_\lambda^{(\alpha)}$ satisfying*

$$(2.3.1) \quad P_\lambda^{(\alpha)} = m_\lambda + \sum_{\mu < \lambda} u_{\lambda\mu}^{(\alpha)} m_\mu \text{ for suitable coefficients } u_{\lambda\mu}^{(\alpha)},$$

$$(2.3.2) \quad \langle P_\lambda^{(\alpha)}, P_\mu^{(\alpha)} \rangle = 0 \text{ if } \lambda \neq \mu.$$

The uniqueness is clear since the basis $\{P_\lambda^{(\alpha)}\}_\lambda$ is obtained by the Gram-Schmidt orthogonalization from $\{m_\lambda\}_\lambda$ with respect to *any* total order compatible with the dominance order. The nontrivial point lies in (2.3.1) where the summation is over μ which is smaller than λ with respect to the dominance order.

Our geometric construction gives a new proof of this theorem.

The “integral form” $J_\lambda^{(\alpha)}$ of $P_\lambda^{(\alpha)}$ is defined by the normalization

$$(2.4) \quad J_\lambda^{(\alpha)} \stackrel{\text{def.}}{=} c_\lambda(\alpha) P_\lambda^{(\alpha)}, \quad c_\lambda(\alpha) \stackrel{\text{def.}}{=} \prod_{s \in \lambda} (\alpha a(s) + l(s) + 1),$$

where “ $s \in \lambda$ ” means that s is a box of the Young diagram corresponds to λ , and $a(s)$ (resp. $l(s)$) is the *arm* (resp. *leg*) length defined by

$$(2.5) \quad \begin{array}{|c|c|c|c|} \hline & & & \\ \hline & \heartsuit & & \\ \hline & \heartsuit & & \\ \hline & s & \spadesuit & \spadesuit \\ \hline & & & \\ \hline \end{array} \quad \begin{array}{l} a(s) = \text{number of } \heartsuit \\ l(s) = \text{number of } \spadesuit \end{array}$$

Since our diagram is rotated, there is no reason to call them arm and leg. But we follow the traditional convention.

The *augmented* monomial symmetric function is defined by

$$\tilde{m}_\lambda \stackrel{\text{def.}}{=} u_\lambda m_\lambda, \quad u_\lambda \stackrel{\text{def.}}{=} \prod_k m_k \quad \text{for } \lambda = (1^{m_1} 2^{m_2} \dots).$$

Macdonald conjectured [17, 10.26?] that $J_\lambda^{(\alpha)}$ is expressed as a linear combination of augmented monomial symmetric function $\tilde{m}_\lambda$ with coefficients in $\mathbb{Z}_{\geq 0}[\alpha]$. This conjecture was proved affirmatively by Knop and Sahi [16] by a combinatorial method.

2(ii). Equivariant Cohomology. We define the equivariant cohomology using the Borel construction. We assume the group is S^1 , though the adaptation to the general compact Lie group is straightforward. Let $ES^1 \rightarrow BS^1$ be the universal S^1 -bundle which is given by the inductive limit of the Hopf fibration $S^{2n+1} \rightarrow \mathbb{CP}^n$. For a topological space M with a circle action, let $M_{S^1} = ES^1 \times_{S^1} M$. We have a projection $M_{S^1} \rightarrow BS^1$, which is a fibration with fiber M . Then, the equivariant cohomology $H_{S^1}^*(M)$ is, by definition, the cohomology of M_{S^1} . Similarly, the equivariant cohomology with compact support, denoted by $H_{S^1, c}^*(M)$ is defined by the cohomology of M_{S^1} with compact support in the fiber direction of $M_{S^1} \rightarrow BS^1$. We assume coefficients are complex numbers.

If M is the space pt consisting of a single point with a trivial S^1 -action, we have $H_{S^1}^*(pt) = H^*(BS^1)$. Since BS^1 is the infinite dimensional projective space $\mathbb{CP}^\infty$, $H^*(BS^1)$ is the polynomial ring with a generator u in $H^2(BS^1)$. We normalize u to be the first Chern class of the tautological line bundle, i.e., the dual of the hyperplane bundle.

By the projection $M_{S^1} \rightarrow BS^1$, we have $H_{S^1}^*(pt) \cong \mathbb{C}[u]$ -module structures on $H_{S^1}^*(M)$ and $H_{S^1, c}^*(M)$.

In order to relate the equivariant cohomology $H_{S^1}^*(M)$ to the ordinary cohomology $H^*(M)$, we consider the Leray-Serre spectral sequence associated with the fibration $M_{S^1} \rightarrow BS^1$. It is a spectral sequence converging to $H^*(M_{S^1}) = H_{S^1}^*(M)$ with E_2 -term $E_2^{p,q} = H^p(BS^1) \otimes H^q(M)$. (Note $\pi_1(BS^1) = 0$.) The (decreasing) filtration is given by

$$F^p H_{S^1}^*(M) = \{u^p \varphi \mid \varphi \in H_{S^1}^*(M)\}.$$

Then Kirwan proved

Theorem 2.6. [15, 5.8] *Let M be a compact symplectic manifold with a Hamiltonian S^1 -action. Then the Leray-Serre spectral sequence associated with $M_{S^1} \rightarrow BS^1$ degenerates at the E_2 -term. Thus we have*

$$F^p H_{S^1}^{p+q}(M) / F^{p+1} H_{S^1}^{p+q}(M) \cong H^p(BS^1) \otimes H^q(M).$$

Kirwan's proof works for a noncompact symplectic manifold provided $f^{-1}((-\infty, c])$ is compact for all $c \in \mathbb{R}$, where f is the moment map associated with the Hamiltonian S^1 -action (see also [19, Chapter 5]). Unfortunately, we do not have a natural Kähler metric on the Hilbert scheme of points, so we could not check the moment map satisfies the above condition. However, we can take another route to save Kirwan's argument. In stead of the gradient flow of the moment map, we use the $\mathbb{C}^*$ -action which extends the S^1 -action. (See [19, Chapter 7].)

We have another short-cut when we assume X is the total space of a line bundle over $\mathbb{CP}^1$. Since the ordinary cohomology groups vanish in odd degree, it is obvious that the spectral sequence degenerates at E_2 -term.

3. HOMOLOGY GROUP OF THE HILBERT SCHEME

The purpose of this section is to identify the complexified ring of symmetric functions $\Lambda \otimes \mathbb{C}$ with the homology group of the Hilbert scheme of X . In fact, the result of this section holds when X is the total space of a line bundle L over *any* compact Riemann surface C .

Let X be as above. It is a nonsingular 2-dimensional quasi-projective surface containing C as the 0-section.

Let $X^{[n]}$ be the Hilbert scheme parameterizing 0-dimensional subschemes of length n . By a result of Fogarty [8], it is a nonsingular $2n$ -dimensional variety.

Let us consider the homology group $H_*(X^{[n]})$ of the Hilbert scheme. In [18], we have constructed an action of the Heisenberg algebra on the direct sum $\bigoplus_n H_*(X^{[n]})$. Let us briefly recall the construction. For each $i \in \mathbb{Z} \setminus \{0\}$, let $P_C[i]$ be a subvariety consisting $(\mathcal{J}_1, \mathcal{J}_2) \in \coprod_n X^{[n-i]} \times X^{[n]}$ such that

$$(3.1) \quad \begin{cases} \mathcal{J}_1 \supset \mathcal{J}_2, \text{ Supp}(\mathcal{J}_1/\mathcal{J}_2) = \{x\} \text{ for some } x \in C & (\text{when } i > 0) \\ \mathcal{J}_1 \subset \mathcal{J}_2, \text{ Supp}(\mathcal{J}_2/\mathcal{J}_1) = \{x\} \text{ for some } x \in C & (\text{when } i < 0) \end{cases},$$

where we consider points in $X^{[n]}$ as ideals of $\mathcal{O}_X$. Let p_a be the projection to the a th factor in the product $X^{[n-i]} \times X^{[n]}$. Note that the restriction of the projection $p_2: P_C[i] \rightarrow X^{[n]}$ is proper. We define a homomorphism $H_*(X^{[n]}) \rightarrow H_*(X^{[n-i]})$ by

$$\varphi \longmapsto p_{1*} (p_2^* \varphi \cap [P_C[i]]),$$

where $p_2^* \varphi \cap$ means the cap product of the pull-back of the Poincaré dual of φ by p_2^* . Since $p_2: P_C[i] \rightarrow X^{[n]}$ is proper, the support of $p_2^* \varphi \cap [P_C[i]]$ is compact, and hence $p_{1*} (p_2^* \varphi \cap [P_C[i]])$ can be defined. Moving n , we get an endomorphism on $\bigoplus_n H_*(X^{[n]})$, which we denote by the same symbol $P_C[i]$ for brevity. Since $P_C[i] \cap (X^{[n-i]} \times X^{[n]})$ is a $(2n - i)$ -dimensional subvariety (see [18, §3] or [19, §8.3] for the proof), $P_C[i]$ maps $H_{2n+k}(X^{[n]})$ to $H_{2(n-i)+k}(X^{[n-i]})$. In particular, the middle degree part ($k = 0$) is preserved. Then the main result of [18] is the following commutator relation

$$(3.2) \quad [P_C[i], P_C[j]] = (-1)^{i-1} i \delta_{i+j,0} \langle C, C \rangle \text{id}.$$

(The factor $(-1)^{i-1} i$, which was not determined in [18], was given by Ellingsrud-Strømme [7]. See also [19, Chapter 9] for a proof in the spirit of this paper.)

Let 1 be the generator of $H_0(X^{[0]}) = \mathbb{C}$. Applying $P_C[i]$ successively on 1, we get a subspace in $\bigoplus_n H_{2n}(X^{[n]})$ which is the irreducible representation of the Heisenberg algebra.

For each partition $\lambda = (1^{m_1} 2^{m_2} \dots)$ define

$$P^\lambda C \stackrel{\text{def.}}{=} P_C[-1]^{m_1} P_C[-2]^{m_2} \dots 1 \in H_{2|\lambda|}(X^{[|\lambda|]}).$$

Since the representation of the Heisenberg algebra generated by $P_C[i]$'s is irreducible, $P^\lambda C$'s are linearly independent.

On the other hand, by the formula of Göttsche [10, 12] for the Poincaré polynomial $P_t(X^{[n]})$ of $X^{[n]}$ (see also [19, Chapter 7] for the proof based on Morse theory) we have

$$\sum_{n=0}^{\infty} q^n P_t(X^{[n]}) = \prod_{m=1}^{\infty} \frac{(1 + t^{2m-1} q^m)^{b_1(X)}}{(1 - t^{2m-2} q^m)(1 - t^{2m} q^m)}.$$

Hence

$$(3.3) \quad \sum_{n=0}^{\infty} q^n \dim H_{2n}(X^{[n]}) = \prod_{m=1}^{\infty} \frac{1}{1 - q^m}.$$

The right hand side of (3.3) is the same as the character of the irreducible representation of the Heisenberg algebra, hence $P^\lambda C$'s span $\bigoplus_n H_{2n}(X^{[n]})$. In other words, $\{P^\lambda C\}_\lambda$ is a basis for $\bigoplus_n H_{2n}(X^{[n]})$.

We identify $\bigoplus_n H_{2n}(X^{[n]})$ with the polynomial ring $\mathbb{C}[p_1, p_2, \dots]$ (and hence with the complexified ring of symmetric functions $\Lambda \otimes \mathbb{C}$) by

$$(3.4) \quad p_\lambda = p_1^{m_1} p_2^{m_2} \dots \longmapsto P^\lambda C = P_C[-1]^{m_1} P_C[-2]^{m_2} \dots 1 \quad \text{for } \lambda = (1^{m_1} 2^{m_2} \dots).$$

Then the operator $P_C[-i]$ corresponds to the multiplication by p_i when $i > 0$. The main result of this section is

Theorem 3.5. (1) *The direct sum $\bigoplus_n H_{2n}(X^{[n]})$ of the middle degree homology group is isomorphic to the complexified ring of symmetric functions $\Lambda \otimes \mathbb{C}$ under the identification (3.4).*

(2) *The intersection pairing $\langle \cdot, \cdot \rangle$ on $H_{2n}(X^{[n]})$ is given by*

$$\langle P^\lambda C, P^\mu C \rangle = (-1)^n \delta_{\lambda\mu} z_\lambda (-\langle C, C \rangle)^{l(\lambda)},$$

where λ, μ is a partition of n .

If we define a new inner product by $\langle \cdot | \cdot \rangle \stackrel{\text{def.}}{=} (-1)^n \langle \cdot, \cdot \rangle$ on $H_{2n}(X^{[n]})$, it is equal to the one used for the definition of Jack's symmetric functions, where the parameter α is $-\langle C, C \rangle$.

Proof of Theorem 3.5. The only remaining is to prove the statement (2).

We identify $\bigoplus_n H_{2n}(X^{[n]})$ with $\mathbb{C}[p_1, p_2, \dots]$ by (1). Then the commutation relation (3.2) means that the operator $P_C[i]$ for $i > 0$ corresponds to

$$(-1)^{i-1} i \langle C, C \rangle \frac{\partial}{\partial p_i}.$$

Hence for $i, j > 0$, we have

$$\begin{aligned} [P_C[i]^m, P_C[-j]^n] &= \left((-1)^{i-1} i \langle C, C \rangle \frac{\partial}{\partial p_i} \right)^m p_j^n \\ (3.6) \quad &= \begin{cases} n(n-1) \cdots (n-m+1) \delta_{ij} \{ (-1)^{i-1} i \langle C, C \rangle \}^m P_C[-j]^{n-m}, & \text{for } n \geq m \\ 0, & \text{for } n < m. \end{cases} \end{aligned}$$

By construction, $P_C[i]$ is the adjoint of $P_C[-i]$ with respect to the intersection form. For $\lambda = (1^{m_1} 2^{m_2} \dots)$, $\mu = (1^{n_1} 2^{n_2} \dots)$ let $\bar{\lambda} = (2^{m_2} 3^{m_3} \dots)$, $\bar{\mu} = (2^{n_2} 3^{n_3} \dots)$. Then we have

$$\begin{aligned} \langle P^\lambda C, P^\mu C \rangle &= \langle P_C[-1]^{m_1} P^{\bar{\lambda}} C, P_C[-1]^{n_1} P^{\bar{\mu}} C \rangle \\ &= \langle P_C[1]^{n_1} P_C[-1]^{m_1} P^{\bar{\lambda}} C, P^{\bar{\mu}} C \rangle. \end{aligned}$$

Since $P_C[-1]$ commutes with $P_C[i]$ for $i \neq -1$, we have

$$\begin{aligned} P_C[1]^{n_1} P_C[-1]^{m_1} P^{\bar{\lambda}} C &= [P_C[1]^{n_1}, P_C[-1]^{m_1}] P^{\bar{\lambda}} C \\ &= \begin{cases} m_1! \langle C, C \rangle^{m_1} P^{\bar{\lambda}} C & \text{for } m_1 = n_1 \\ 0, & \text{otherwise,} \end{cases} \end{aligned}$$

where we have used (3.6). Hence we have

$$\langle P^\lambda C, P^\mu C \rangle = \delta_{m_1, n_1} m_1! \langle C, C \rangle^{m_1} \langle P^{\bar{\lambda}} C, P^{\bar{\mu}} C \rangle.$$

Inductively, we get

$$\begin{aligned} \langle P^\lambda C, P^\mu C \rangle &= \delta_{\lambda\mu} \prod_i ((-1)^{i-1} i \langle C, C \rangle)^{m_i} m_i! \\ &= \delta_{\lambda\mu} z_\lambda (-1)^{\sum i m_i} (-\langle C, C \rangle)^{\sum m_i}. \end{aligned}$$

Since $\sum i m_i = n$, $\sum m_i = l(\lambda)$, we get the assertion. $\square$

4. CERTAIN SUBVARIETIES AND MONOMIAL SYMMETRIC FUNCTIONS

In this section we define certain middle dimensional subvarieties parametrized by partitions and identify them with monomial symmetric functions under the isomorphism given in Theorem 3.5. The subvarieties were first introduced by Grojnowski [13], and their identification with monomial symmetric functions was proved in [19, Chapter 9]. We reproduce them for the sake of the reader.

Let X be as in §3, i.e., the total space of a line bundle over a compact Riemann surface C . Let us consider the $\mathbb{C}^*$ -action on X given by the multiplication on fibers. It induces an action on the Hilbert scheme $X^{[n]}$. Let us consider a subvariety of the Hilbert scheme $X^{[n]}$ defined by

$$(4.1) \quad \{Z \in X^{[n]} \mid \text{Supp}(\mathcal{O}_Z) \subset C\}.$$

Since the $\mathbb{C}^*$ -action retracts $X^{[n]}$ to a neighborhood of the above subvariety, $X^{[n]}$ is homotopically equivalent to the subvariety. It was pointed out by Grojnowski [13] (see [19, Chapter 7] for the proof) that its irreducible components $L^\lambda C$ are indexed by a partition λ as follows: first let $\pi: X^{[n]} \rightarrow S^n X$ be the Hilbert-Chow morphism which assigns to a closed subscheme Z of X , the 0-cycle consisting of the points of Z with multiplicities given by the length of the local rings on Z . Then (4.1) is given by $\pi^{-1}(S^n C)$, where $S^n C$ is considered as a subvariety of $S^n X$. We have a stratification of $S^n C$ given by

$$S^n C = \bigcup_{\lambda} S_{\lambda}^n C, \quad \text{where } S_{\lambda}^n C \stackrel{\text{def.}}{=} \left\{ \sum_{i=1}^k \lambda_i [x_i] \in S^n C \mid x_i \neq x_j \text{ for } i \neq j \right\},$$

where $\lambda = (\lambda_1, \lambda_2, \dots)$ runs over partitions of n . Now consider a locally closed subvariety $\pi^{-1}(S_{\lambda}^n C)$ of (4.1). By a result of Briançon [4], the Hilbert-Chow morphism π is semismall (see also [12] or [19, Chapter 6]), the dimension of the fiber of π over a point in $S_{\lambda}^n C$ is $n - l(\lambda)$. Hence the dimension of $\pi^{-1}(S_{\lambda}^n C)$ is equal to n , which is independent of the partition λ . Moreover, the fiber of π is irreducible again by a result of Briançon [4]. Thus the irreducible components of (4.1) are given by

$$(4.2) \quad L^{\lambda} C \stackrel{\text{def.}}{=} \text{Closure of } \pi^{-1}(S_{\lambda}^n C).$$

We need another definition of $L^{\lambda} C$. Let us consider a fixed point $\mathcal{J}$ of the $\mathbb{C}^*$ -action in $X^{[n]}$. Since the fixed point in X is the zero section C , we have $\text{Supp}(\mathcal{O}_X/\mathcal{J}) \subset C$. Let us decompose $\mathcal{J}$ as $\mathcal{J}_1 \cap \dots \cap \mathcal{J}_m$ according to the support, i.e., $\text{Supp}(\mathcal{O}_X/\mathcal{J}_k) = \{x_k\}$ and $x_k \neq x_l$ for $k \neq l$. For each k , we take a coordinate system (z, ξ) around x_k where z is a coordinate of C around x_k , and ξ is a fiber coordinate of L_{x_k} . Then $\mathcal{J}_k$ is generated by monomials in z and ξ :

$$\mathcal{J}_k = (\xi^{\lambda_1^{(k)}}, z\xi^{\lambda_2^{(k)}}, \dots, z^{N-1}\xi^{\lambda_N^{(k)}}, z^N),$$

for some partition $\lambda^{(k)} = (\lambda_1^{(k)}, \lambda_2^{(k)}, \dots)$ with $N = l(\lambda^{(k)})$. For a partition λ and a point $x \in C$, let

$$(4.3) \quad \mathcal{J}_{\lambda, x} \stackrel{\text{def.}}{=} (\xi^{\lambda_1}, z\xi^{\lambda_2}, \dots, z^{N-1}\xi^{\lambda_N}, z^N),$$

where $N = l(\lambda)$ and (z, ξ) is the coordinate system around x . (See Figure 6.1 for the visualization of $\mathcal{J}_{\lambda, x}$.) Thus the fixed point can be written as $\mathcal{J} = \bigcap \mathcal{J}_{\lambda^{(k)}, x_k}$ for some distinct points x_k 's in C and partitions $\lambda^{(k)}$.

If x_k approaches to x_l , $\mathcal{J}_{\lambda^{(k)}, x_k} \cap \mathcal{J}_{\lambda^{(l)}, x_l}$ converges to $\mathcal{J}_{\lambda^{(k)} \cup \lambda^{(l)}, x_l}$. This shows that the fixed point components are parametrized by partitions $\lambda = \bigcup \lambda^{(k)}$. Let us denote by $S^\lambda C$ the corresponding component. By the above discussion, we have

$$S^\lambda C \stackrel{\text{def.}}{=} \left\{ \mathcal{J} = \bigcap \mathcal{J}_{\lambda^{(k)}, x_k} \mid x_i \in C, x_i \neq x_j \text{ for } i \neq j, \lambda = \bigcup \lambda^{(k)} \right\}.$$

As a complex manifold, $S^\lambda C$ is isomorphic to a product of symmetric product

$$S^\lambda C \cong S^{m_1} C \times S^{m_2} C \times \cdots,$$

where $\lambda = (1^{m_1} 2^{m_2} \dots)$. Now another description of $L^\lambda C$ is

$$(4.4) \quad L^\lambda C = \text{Closure of } \{ \mathcal{J} \in X^{[n]} \mid \lim_{t \rightarrow \infty} t \cdot \mathcal{J} \in S^\lambda C \},$$

where $t \cdot$ denotes the $\mathbb{C}^*$ -action. In fact, since the Hilbert-Chow morphism $\pi: X^{[n]} \rightarrow S^n X$ is $\mathbb{C}^*$ -equivariant, $t \cdot \mathcal{J}$ stays a compact set, or equivalently converges to a fixed point if and only if $\text{Supp}(\mathcal{O}_X/\mathcal{J}) \subset C$. Moreover, $\lim_{t \rightarrow \infty} t \cdot \mathcal{J}$ is contained in the open stratum

$$\{ \mathcal{J}_{(\lambda_1), x_1} \cap \cdots \cap \mathcal{J}_{(\lambda_N), x_N} \mid x_i \in C, x_i \neq x_j \text{ for } i \neq j \}$$

of $S^\lambda C$ if and only if $\mathcal{J} \in \pi^{-1}(S^\lambda C)$. This shows the identification (4.4).

Remark 4.5. As is presented in [19, Chapter 7], we can start from (4.4) as a definition of $L^\lambda C$. Studying the weight decomposition of the tangent space at a point in $S^\lambda C$, we can prove that (4.4) is n -dimensional (cf. discussion for Proposition 6.5). The irreducibility of (4.4) follows from that of $S^\lambda C$. Hence (4.4) is an irreducible component of (4.1).

Theorem 4.6. *Under the isomorphism $\Lambda \otimes \mathbb{C} \cong \bigoplus H_{2n}(X^{[n]})$ given in Theorem 3.5, the class $[L^\lambda C]$ corresponds to the monomial symmetric function m_λ .*

Proof. We shall show

$$(4.7) \quad P_C[-i][L^\lambda C] = \sum_{\mu} a_{\lambda\mu} [L^\mu C], \quad \text{for any } i \in \mathbb{Z}_{>0},$$

where the summation is over partitions μ of $|\lambda| + i$ which is obtained as follows:

- (4.8.1) Add i to a term in λ , say λ_k (possibly 0).
- (4.8.2) Then arrange it in descending order.
- (4.8.3) Define the coefficient $a_{\lambda\mu}$ by $\#\{l \mid \mu_l = \lambda_k + i\}$.

This is the same as the relation between the power sum and the monomial symmetric function:

$$p_i m_\lambda = \sum_{\mu} a_{\lambda\mu} m_\mu.$$

By induction, we have $[L^\lambda C] = m_\lambda$ under the identification $P_C[-i] = p_i$.

The equation (4.7) is proved by studying the intersection product. Let $p_a: X^{[n]} \times X^{[n]} \rightarrow X^{[n]}$ be the projection onto the a th factor ($a = 1, 2$). By the definition (3.1), we represent $P_C[-i]$ as a subvariety. Then its set theoretical intersection with $p_2^{-1}(L^\lambda C)$ is

$$(4.9) \quad \{(\mathcal{J}_1, \mathcal{J}_2) \mid \mathcal{J}_2 \in L^\lambda C, \mathcal{J}_1 \subset \mathcal{J}_2, \text{Supp}(\mathcal{J}_2/\mathcal{J}_1) = \{x\} \text{ for some } x \in C\}.$$

Let $\mu = (\mu_1, \mu_2, \dots)$ be a partition of $|\lambda| + i$ which does not necessarily satisfy the condition in (4.8). Let $N = l(\mu)$. Since $\{[L^\mu C]\}_\mu$ is a basis for $H_{2|\mu|}(X^{[|\mu|]})$, the left hand side of (4.7) can be written as a linear combination of $[L^\mu C]$'s. In order to determine the coefficients of $[L^\mu C]$ in $P_C[-i][L^\lambda C]$, it is enough to take arbitrary point $\mathcal{J}_1$ in $L^\mu C$ and restrict cycles to a neighborhood of $\mathcal{J}_1$. We choose the point $\mathcal{J}_1 = \mathcal{J}_{(\mu_1), x_1} \cap \dots \cap \mathcal{J}_{(\mu_N), x_N} \in L^\mu C$ where x_k 's are distinct points in C . Here $\mathcal{J}_{(\mu_k), x_k} = (\xi^{\mu_k}, z)$ for the coordinate (z, ξ) around x_k . (See (4.3) for the definition of $\mathcal{J}_{\lambda, x}$.)

Suppose this point $\mathcal{J}_1$ is contained in the image of (4.9) under the projection p_1 , i.e., there exists $\mathcal{J}_2$ such that $(\mathcal{J}_1, \mathcal{J}_2)$ is a point in (4.9). Then the point x must be one of x_k 's, and

$$\mathcal{J}_2 = \mathcal{J}_{(\mu_1), x_1} \cap \dots \cap \mathcal{J}_{(\mu_{k-1}), x_{k-1}} \cap \mathcal{J}_{(\mu_k - i), x_k} \cap \mathcal{J}_{(\mu_{k+1}), x_{k+1}} \cap \dots \cap \mathcal{J}_{(\mu_N), x_N},$$

i.e., $\mathcal{J}_2$ is obtained from $\mathcal{J}_1$ by replacing $\mathcal{J}_{(\mu_k), x_k}$ by $\mathcal{J}_{(\mu_k - i), x_k}$. Since $\mathcal{J}_2$ must be a point in $L^\lambda C$, μ is obtained by (a) adding i to λ_k , and then (b) arranging in descending order.

Moreover, if $a_{\lambda\mu}$ is as in (4.8), there are exactly $a_{\lambda\mu}$ choices of x_k 's. This explains the coefficient $a_{\lambda\mu}$ in the formula. Thus the only remaining thing to check is that each choice of $(\mathcal{J}_1, \mathcal{J}_2)$ contributes to $P_C[-i][L^\lambda C]$ by $[L^\mu C]$. This will be shown by checking

(4.10.1) $P_C[-i]$ and $p_2^{-1}(L^\lambda C)$ intersect transversally,

(4.10.2) the intersection (4.9) is isomorphic to $L^\mu C$ under the first projection p_1 , in a neighborhood of $(\mathcal{J}_1, \mathcal{J}_2)$.

Since $\mathcal{J}_1$ and $\mathcal{J}_2$ are isomorphic outside x_k , we can restrict our concern to $\mathcal{J}_{(\mu_k), x_k}$ and $\mathcal{J}_{(\mu_k - i), x_k}$. We take the following coordinate neighborhood around $(\mathcal{J}_{(\mu_k), x_k}, \mathcal{J}_{(\mu_k - i), x_k})$ in $X^{[\mu_k]} \times X^{[\mu_k - i]}$:

$$\begin{aligned} & \{((\xi^{\mu_k} + f_1(\xi), z + g_1(\xi)), (\xi^{\mu_k - i} + f_2(\xi), z + g_2(\xi))) \mid f_1, g_1, f_2, g_2 \text{ as follows}\} \\ & f_1(\xi) = a_1 \xi^{\mu_k - 1} + a_2 \xi^{\mu_k - 2} + \dots + a_{\mu_k}, \\ & g_1(\xi) = b_1 + b_2 \xi + \dots + b_{\mu_k - i} \xi^{\mu_k - i - 1} \\ & \quad + (b_{\mu_k - i + 1} + b_{\mu_k - i + 2} \xi + \dots + b_{\mu_k} \xi^{i - 1})(\xi^{\mu_k - i} + f_2(\xi)) \\ & f_2(\xi) = a'_1 \xi^{\mu_k - i - 1} + a'_2 \xi^{\mu_k - i - 2} + \dots + a'_{\mu_k - i}, \\ & g_2(\xi) = b'_1 + b'_2 \xi + \dots + b'_{\mu_k - i} \xi^{\mu_k - i - 1} \end{aligned}$$

where $(a_1, \dots, a_{\mu_k}, b_1, \dots, b_{\mu_k})$ (resp. $(a'_1, \dots, a'_{\mu_k - i}, b'_1, \dots, b'_{\mu_k - i})$) is in a neighborhood of 0 in $\mathbb{C}^{2\mu_k}$ (resp. $\mathbb{C}^{2(\mu_k - i)}$). Then the above ideal is contained in $P_C[-i]$ if and only if the followings hold

$$(4.11.1) \quad \xi^{\mu_k} + f_1(\xi) = \xi^i (\xi^{\mu_k - i} + f_2(\xi)),$$

$$(4.11.2) \quad g_1(\xi) - g_2(\xi) \text{ is divisible by } \xi^{\mu_k - i} + f_2(\xi).$$

Namely, the defining equation for $P_C[-i]$ is

$$\begin{aligned} a_1 &= a'_1, a_2 = a'_2, \dots, a_{\mu_k-i} = a'_{\mu_k-i}, \\ a_{\mu_k-i+1} &= \dots = a_{\mu_k} = 0, \\ b_1 &= b'_1, b_2 = b'_2, \dots, b_{\mu_k-i} = b'_{\mu_k-i}. \end{aligned}$$

On the other hand, the defining equation for $p_2^{-1}(L^\lambda C)$ is

$$a'_1 = a'_2 = \dots = a'_{\mu_k-i} = 0.$$

Now our assertions (4.10) are immediate. $\square$

Our next task is to explain a geometric meaning of the dominance order (2.2). It is given by modifying the stratification introduced in [4, 14].

For $i \geq 0$, let $\mathcal{O}_X(-iC)$ be the sheaf of functions vanishing to order $\geq i$ along the zero section C . Let $\mathcal{J} \subset \mathcal{O}_X$ be an ideal of colength n such that the support of $\mathcal{O}_X/\mathcal{J}$ is contained in C . We consider the sequence $(\lambda'_1, \lambda'_2, \dots)$ of nonnegative integers given by

$$\lambda'_i(\mathcal{J}) \stackrel{\text{def.}}{=} \text{length} \left(\frac{\mathcal{O}_X(-(i-1)C)}{\mathcal{J} \cap \mathcal{O}_X(-(i-1)C) + \mathcal{O}_X(-iC)} \right).$$

The reason why we put the prime become clear later. The sequence in [4, 14] was defined by replacing $\mathcal{O}_X(-iC)$ by $\mathfrak{m}_x^i$ where $\mathfrak{m}_x$ is the maximal ideal corresponding to a point x . As in [14, Lemma 1.1], we have $\mathcal{J} \supset \mathcal{O}_X(-nC)$, hence $\lambda'_i(\mathcal{J}) = 0$ for $i \geq n+1$. From the exact sequence

$$0 \rightarrow \frac{\mathcal{O}_X(-iC)}{\mathcal{J} \cap \mathcal{O}_X(-iC)} \rightarrow \frac{\mathcal{O}_X(-(i-1)C)}{\mathcal{J} \cap \mathcal{O}_X(-(i-1)C)} \rightarrow \frac{\mathcal{O}_X(-(i-1)C)}{\mathcal{J} \cap \mathcal{O}_X(-(i-1)C) + \mathcal{O}_X(-iC)} \rightarrow 0,$$

we have

$$\sum_{i=1}^n \lambda'_i(\mathcal{J}) = n.$$

Let us decompose the ideal $\mathcal{J}$ by its support, i.e., $\mathcal{J} = \mathcal{J}_1 \cap \dots \cap \mathcal{J}_N$ such that $\{\text{Supp}(\mathcal{O}_X/\mathcal{J}_k)\}_k$ are N distinct points. By definition, we have

$$(4.12) \quad \lambda'_i(\mathcal{J}) = \sum_{k=1}^N \lambda'_i(\mathcal{J}_k).$$

Suppose that $\mathcal{J}$ satisfies $\text{Supp}(\mathcal{O}_X/\mathcal{J}) = \{x\}$ for some $x \in C$. We take a coordinate system (z, ξ) around x where ξ is a coordinate for the fiber. If $\xi^i f_1(z), \dots, \xi^i f_d(z)$ form a basis of

$$\frac{\mathcal{O}_X(-iC)}{\mathcal{J} \cap \mathcal{O}_X(-iC) + \mathcal{O}_X(-(i+1)C)},$$

Then $\xi^{i-1}f_1(z), \dots, \xi^{i-1}f_d(z)$ are linearly independent in

$$\frac{\mathcal{O}_X(-(i-1)C)}{\mathcal{J} \cap \mathcal{O}_X(-(i-1)C) + \mathcal{O}_X(-iC)}.$$

Hence we have $\lambda'_i(\mathcal{J}) \geq \lambda'_{i+1}(\mathcal{J})$. Thus $(\lambda'_1(\mathcal{J}), \lambda'_2(\mathcal{J}), \dots)$ is a partition of n . By (4.12), the same is true for general $\mathcal{J}$ which do not necessarily satisfy $\text{Supp}(\mathcal{O}_X/\mathcal{J}) = \{x\}$. Let us denote the partition by $\lambda'(\mathcal{J})$.

For a partition $\lambda' = (\lambda'_1, \lambda'_2, \dots)$ of n , let $W^{\lambda'}$ be the set of ideals $\mathcal{J} \subset \mathcal{O}_X$ with colength $n = |\lambda'|$ such that $\mathcal{O}_X/\mathcal{J}$ is supported on C and $\lambda'(\mathcal{J}) = \lambda'$. Since

$$\text{length}(\mathcal{O}_X(-iC)/\mathcal{J} \cap \mathcal{O}_X(-iC)) \leq \sum_{j=i+1}^n \lambda'_j = n - \sum_{j=1}^i \lambda'_j$$

is a closed condition on $\mathcal{J}$, the union

$$\bigcup_{\mu' \geq \lambda'} W^{\mu'}$$

is a closed subset of $\{\mathcal{J} \in X^{[n]} \mid \text{Supp}(\mathcal{O}_X/\mathcal{J}) \subset C\}$. Thus we have

$$(4.13) \quad \text{Closure of } W^{\lambda'} \subset \bigcup_{\mu' \geq \lambda'} W^{\mu'}.$$

Suppose that λ' is the conjugate of λ as in (2.1). We get the following third description of $L^\lambda C$.

Proposition 4.14. $L^\lambda C = \text{Closure of } W^{\lambda'}$.

Proof. Let us write $\lambda = (\lambda_1, \dots, \lambda_N)$ with $N = l(\lambda)$. By (4.4), a generic point $\mathcal{J}$ in $L^\lambda C$ satisfies $\lim_{t \rightarrow \infty} t \cdot \mathcal{J} = \mathcal{J}_{(\lambda_1), x_1} \cap \dots \cap \mathcal{J}_{(\lambda_N), x_N}$ such that x_i 's are distinct points in C . Since the support of $\mathcal{J}$ cannot move as $t \rightarrow \infty$, we can decompose $\mathcal{J} = \mathcal{J}_1 \cap \dots \cap \mathcal{J}_N$ such that $\text{Supp}(\mathcal{O}_X/\mathcal{J}_k) = \{x_k\}$.

Take a coordinate system (z, ξ) around x_k as before. Then $\mathcal{J}_{(\lambda_k), x_k}$ was defined by (z, ξ^{λ_k}) . Since $\lim_{t \rightarrow \infty} t \cdot \mathcal{J}_k = \mathcal{J}_{(\lambda_k), x_k}$, we have

$$\mathcal{J}_k = (\xi^{\lambda_k}, z + a_1 \xi + a_2 \xi^2 + \dots + a_{\lambda_k-1} \xi^{\lambda_k-1})$$

for some $a_1, \dots, a_{\lambda_k-1}$ (cf. the proof for Theorem 4.6). Then $\lambda'(\mathcal{J}_k) = (1^{\lambda_k})$. By (4.12), we have

$$\lambda'(\mathcal{J}) = \lambda'.$$

This shows $L^\lambda C \subset \text{Closure of } W^{\lambda'}$.

Conversely, suppose $\mathcal{J}$ is a point in $W^{\lambda'}$. Let

$$\mathcal{J}_i \stackrel{\text{def.}}{=} \frac{\mathcal{J} \cap \mathcal{O}_X(-(i-1)C)}{\mathcal{J} \cap \mathcal{O}_X(-iC)}.$$

Then $\mathcal{J}' \stackrel{\text{def.}}{=} \bigoplus \mathcal{J}_i$ satisfies $\lim_{t \rightarrow \infty} t \cdot \mathcal{J} = \mathcal{J}'$ and $\mathcal{J}' \in W^{\lambda'}$. This shows $W^{\lambda'} \subset L^\lambda C$. $\square$

5. EQUIVARIANT COHOMOLOGY OF HILBERT SCHEMES

Let X be the total space of a line bundle over $\mathbb{CP}^1$ and C the zero section. X is the quotient space of $(\mathbb{C}^2 \setminus \{0\}) \times \mathbb{C}$ by the $\mathbb{C}^*$ -action given by

$$(z_0, z_1, \xi) \mapsto (\lambda z_0, \lambda z_1, \lambda^{-\alpha} \xi) \quad \lambda \in \mathbb{C}^*,$$

where $\alpha = -c_1(L)[C] = -\langle C, C \rangle$. We denote by $[(z_0, z_1, \xi)]$ the equivalence class containing (z_0, z_1, ξ) . The projection $X \rightarrow \mathbb{CP}^1$ is given by $[(z_0, z_1, \xi)] \mapsto [z_0 : z_1]$, hence ξ is a coordinate for the fiber.

We consider the S^1 -action on $X^{[n]}$ induced by the action on X defined by

$$(5.1) \quad [(z_0, z_1, \xi)] \mapsto [(z_0, t^{-1}z_1, t^\alpha \xi)] \quad \text{for } t \in S^1.$$

Note that this differs from the S^1 -action studied in the previous section given by $[(z_0, z_1, \xi)] \mapsto [(z_0, z_1, t\xi)]$. We shall study the equivariant cohomology of $X^{[n]}$ with respect to the above S^1 -action. The reason for using this S^1 -action is to identify the normalization factor (2.4) with the equivariant Euler class (see Proposition 6.5).

Let pt be the space consisting of a single point with a trivial S^1 -action. Let us denote the obvious S^1 -equivariant morphism by $p: X^{[n]} \rightarrow pt$. We have an equivariant push-forward p_* . Define a bilinear form $\langle \cdot, \cdot \rangle_{S^1}: H_{S^1, c}^k(X^{[n]}) \otimes H_{S^1, c}^k(X^{[n]}) \rightarrow H_{S^1}^{2(k-2n)}(pt)$ by

$$(5.2) \quad \langle \varphi, \psi \rangle_{S^1} \stackrel{\text{def.}}{=} p_*(\varphi \cup \psi),$$

where $\cup$ is the cup product

$$\cup: H_{S^1, c}^k(X^{[n]}) \otimes H_{S^1, c}^k(X^{[n]}) \rightarrow H_{S^1, c}^{2k}(X^{[n]}).$$

This is trivial unless k is even and $k \geq 2n$. We assume this condition henceforth. In particular, $\langle \cdot, \cdot \rangle_{S^1}$ is symmetric.

Let $j^*: H_{S^1, c}^k(X^{[n]}) \rightarrow H_{S^1}^k(X^{[n]})$ be the natural homomorphism. Then the cup product $\cup$ factor through j^* :

$$\varphi \cup \psi = \varphi \cup j^* \psi,$$

where $\cup$ in the right hand side is the cup product

$$\cup: H_{S^1, c}^k(X^{[n]}) \otimes H_{S^1}^k(X^{[n]}) \rightarrow H_{S^1, c}^{2k}(X^{[n]}).$$

In particular, $\langle \cdot, \cdot \rangle_{S^1}$ is well-defined on

$$H_{S^1, c}^k(X^{[n]}) / \text{Ker } j^* \cong \text{Im } j^*.$$

Let $X_{S^1}^{[n]} \rightarrow BS^1$ be the fibration used in the definition of the equivariant cohomology. By Theorem 2.6 (see also the discussion after Theorem 2.6), the spectral sequence of the fibration in cohomology degenerates. Let us denote by $F^* H_{S^1}^k(X^{[n]})$ the corresponding decreasing filtration.

Since $H^q(X^{[n]}) = 0$ for $q > 2n$, we have

$$\begin{aligned} F^p H_{S^1}^k(X^{[n]}) &= H_{S^1}^k(X^{[n]}) \quad \text{for } p \leq k - 2n, \\ H_{S^1}^k(X^{[n]}) / F^{k-2n+1} H_{S^1}^k(X^{[n]}) &\cong H_{S^1}^{k-2n}(pt) \otimes H^{2n}(X^{[n]}). \end{aligned}$$

If $j^*\psi \in F^{k-2n+1}H_{S^1}^k(X^{[n]})$, then $\varphi \cup j^*\psi \in F^{2(k-2n)+1}H_{S^1,c}^{2k}(X^{[n]})$. Hence $p_*(\varphi \cup j^*\psi) \in F^{2(k-2n)+1}H_{S^1}^{2(k-2n)}(pt) \cong 0$. In particular, $\langle \cdot, \cdot \rangle_{S^1}$ is well-defined on $\text{Im } j^* / \text{Im } j^* \cap F^{k-2n+1}H_{S^1}^k(X^{[n]})$. Consider the composition

$$(5.3) \quad \begin{aligned} & H_{S^1,c}^k(X^{[n]}) / \text{Rad} \langle \cdot, \cdot \rangle_{S^1} \xrightarrow{j^*} \text{Im } j^* / \text{Im } j^* \cap F^{k-2n+1}H_{S^1}^k(X^{[n]}) \\ & \rightarrow H_{S^1}^k(X^{[n]}) / F^{k-2n+1}H_{S^1}^k(X^{[n]}) \cong H_{S^1}^{k-2n}(pt) \otimes H^{2n}(X^{[n]}), \end{aligned}$$

which is injective by the above discussion.

We have an analogous injective homomorphism between *ordinary* cohomology groups

$$H_c^{2n}(X^{[n]}) / \text{Rad} \langle \cdot, \cdot \rangle \rightarrow H^{2n}(X^{[n]}).$$

However, the bilinear form $\langle \cdot, \cdot \rangle$ on $H_c^{2n}(X^{[n]})$ is nondegenerate by Theorem 3.5(2). Hence $\text{Rad} \langle \cdot, \cdot \rangle = 0$. Moreover, the Poincaré duality implies $\dim H_c^{2n}(X^{[n]}) = \dim H^{2n}(X^{[n]})$. Therefore the natural homomorphism $H_c^{2n}(X^{[n]}) \rightarrow H^{2n}(X^{[n]})$ is an isomorphism.

Theorem 5.4. *Suppose k is even and $k \geq 2n$. Then the composition (5.3)*

$$(5.5) \quad H_{S^1,c}^k(X^{[n]}) / \text{Rad} \langle \cdot, \cdot \rangle_{S^1} \rightarrow H_{S^1}^{k-2n}(pt) \otimes H^{2n}(X^{[n]})$$

is an isomorphism. Moreover, the bilinear form $\langle \cdot, \cdot \rangle_{S^1}$ on the left hand side is equal to one on the right hand side induced by the intersection pairing on $H^{2n}(X^{[n]}) \cong H_c^{2n}(X^{[n]})$.

Proof. The subvariety $L^\lambda C$ defined in (4.2) is invariant under the S^1 -action. Hence its Poincaré dual defines an element in the equivariant cohomology $H_{S^1,c}^{2n}(X^{[n]})$, which we denote by $[L^\lambda C]_{S^1}$. Under (5.3) with $k = 2n$, $[L^\lambda C]_{S^1} \bmod \text{Rad} \langle \cdot, \cdot \rangle_{S^1}$ is mapped to $1 \otimes [L^\lambda C]$. More generally, if u is the generator of $H_{S^1}^*(pt)$ which lives in $H_{S^1}^2(pt)$, $u^{k/2-n}[L^\lambda C]_{S^1}$ is mapped to $u^{k/2-n} \otimes [L^\lambda C]$. This shows that (5.5) is surjective. As we have already seen the injectivity, it is an isomorphism.

Next consider the bilinear forms on the both hand sides of (5.5). The doubt arises from that $H_{S^1}^k(X^{[n]}) \cong \bigoplus_{p+q=k} H_{S^1}^p(pt) \otimes H^q(X^{[n]})$ is not an *algebra* isomorphism, it is only an isomorphism between *vector spaces*. If we consider the graded space $GH_{S^1}^k(X^{[n]})$ instead of $H_{S^1}^k(X^{[n]})$, it becomes an algebra isomorphism. However, as we pointed out above, $F^{k-2n+1}H_{S^1}^k(X^{[n]})$ does not contribute to $\langle \cdot, \cdot \rangle_{S^1}$. Hence we have the assertion. $\square$

6. LOCALIZATION

We assume X is the total space of a line bundle over $\mathbb{CP}^1$ with $\alpha = -c_1(L)[\mathbb{CP}^1] > 0$. Let us consider the S^1 -action given in (5.1). The fixed point set consists of two components

$$\{(1, 0, 0)\} \text{ and } \{(0, 1, \xi) \mid \xi \in \mathbb{C}\}.$$

The first is an isolated point and the latter is isomorphic to $\mathbb{C}$. If $Z \in X^{[n]}$ is fixed by the induced action, its support must be contained in the above fixed point set.

There are a family of fixed points Z_λ with support $[(1, 0, 0)]$ indexed by a partition $\lambda = (\lambda_1, \lambda_2, \dots)$ of n as follows: let (z, ξ) be a coordinate system around $[(1, 0, 0)]$ given

by $(z, \xi) \leftrightarrow [(1, z, \xi)]$. Then the corresponding ideal $\mathcal{J}_\lambda$ is given by

$$\mathcal{J}_\lambda \stackrel{\text{def.}}{=} (\xi^{\lambda_1}, z\xi^{\lambda_2}, \dots, z^{N-1}\xi^{\lambda_N}, z^N),$$

where $N = l(\lambda)$. Namely, $\mathcal{J}_\lambda = \mathcal{J}_{\lambda, [(1,0,0)]}$ in the notation (4.3). The ideal can be visualized by the Young diagram as follows. (Remind that our diagram is rotated by $\pi/2$ from one used in [17].) If we write the monomial $z^{p-1}\xi^{q-1}$ at the position (p, q) , the ideal is generated by monomials outside the Young diagram. For example, the Young diagram in figure 6.1 corresponds to the ideal $(\xi^4, z\xi^3, z^2\xi, z^3)$. Although it is not necessary for our purpose, one can show that these ideals are a complete list of fixed points in $X^{[n]}$ whose support are $[(1, 0, 0)]$ when $\alpha > 0$.




FIGURE 6.1. Young diagram and an ideal

Let $i_\lambda: Z_\lambda \rightarrow X^{[n]}$ be the inclusion. We have a homomorphism between equivariant cohomology groups:

$$H_{S^1}^*(Z_\lambda) \ni \varphi \longmapsto i_{\lambda*}\varphi \in H_{S^1,c}^*(X^{[n]}).$$

Let $e(T_{Z_\lambda}X^{[n]})$ be the S^1 -equivariant Euler class of the tangent space at Z_λ , considered as an element of $H_{S^1}^{4n}(Z_\lambda)$. Then we have

Theorem 6.1. *Let $\langle \cdot, \cdot \rangle_{S^1}$ be the bilinear pairing defined in (5.2). Then we have*

$$\langle i_{\lambda*}\varphi, i_{\mu*}\psi \rangle_{S^1} = \begin{cases} 0 & \text{if } \lambda \neq \mu, \\ p_{\lambda*}(\varphi \cup \psi \cup e(T_{Z_\lambda}X^{[n]})) & \text{if } \lambda = \mu, \end{cases}$$

where $p_\lambda: Z_\lambda \rightarrow pt$ is the obvious map.

Proof. By the projection formula, we have

$$i_{\lambda*}\varphi \cup i_{\mu*}\psi = i_{\mu*}(i_\mu^*i_{\lambda*}\varphi \cup \psi).$$

The homomorphism $i_\mu^*i_{\lambda*}$ is zero when $\lambda \neq \mu$, and is given by the multiplication by the equivariant Euler class $e(T_{Z_\lambda}X^{[n]})$ if $\lambda = \mu$. Therefore,

$$\begin{aligned} \langle i_{\lambda*}\varphi, i_{\mu*}\psi \rangle_{S^1} &= p_*(i_{\lambda*}\varphi \cup i_{\mu*}\psi) = p_*i_{\lambda*}(\varphi \cup \psi \cup e(T_{Z_\lambda}X^{[n]})) \\ &= p_{\lambda*}(\varphi \cup \psi \cup e(T_{Z_\lambda}X^{[n]})) \end{aligned} \quad \square$$

Taking a class from $H_{S^1}^*(Z_\lambda)$ for each λ , we can give a set of mutually orthogonal elements in $H_{S^1, c}^*(X^{[n]})$. These are candidates for Jack polynomials. Our remaining task is to choose classes so that two conditions in Theorem 2.3 will be satisfied. Thus we need to know monomial symmetric functions, or their geometric counterparts $[L^\lambda C]_{S^1}$.

In order to study $[L^\lambda C]_{S^1}$, we need to know the S^1 -module structure of its normal bundle at Z_λ . In fact, the formula will become clearer if we consider the two dimensional torus action given by

$$[(z_0, z_1, \xi)] \mapsto [(z_0, t_1 z_1, t_2 \xi)] \quad \text{for } (t_1, t_2) \in T = S^1 \times S^1.$$

The S^1 -action studied in §4 is given by the restriction to $(1, t) \in T$, while the S^1 -action given by (5.1) is the restriction to (t^{-1}, t^α) . The fixed point Z_λ defined above is also fixed by this torus action. The tangent space $T_{Z_\lambda} X^{[n]}$ at Z_λ is a T -module and have a weight decomposition

$$T_{Z_\lambda} X^{[n]} = \bigoplus_{p, q \in \mathbb{Z}} H(p, q),$$

where $H(p, q) = \{v \in T_{Z_\lambda} X^{[n]} \mid (t_1, t_2) \cdot v = t_1^p t_2^q v \text{ for } (t_1, t_2) \in T\}$.

Lemma 6.2. *The character of the tangent space $T_{Z_\lambda} X^{[n]}$ is given by*

$$\sum_{p, q} \dim_{\mathbb{C}} H(p, q) T_1^p T_2^q = \sum_{s \in \lambda} \left(T_1^{l(s)+1} T_2^{-a(s)} + T_1^{-l(s)} T_2^{a(s)+1} \right).$$

Proof. The corresponding formula was proved in [6] (see also [11, 2.2.5] or [19, Proposition 5.5]) when X is replaced by $\mathbb{C}^2$ with the torus action given by

$$(z_1, z_2) \mapsto (t_1 z_1, t_2 z_2) \quad \text{for } (z_1, z_2) \in \mathbb{C}^2, (t_1, t_2) \in T.$$

(The presentation in [6] or [11] is different from the above. After substituting the arm/leg-length to their formula, we get the above formula.) Since the exponential map gives a T -equivariant isomorphism between a neighborhood of $0 \in \mathbb{C}^2$ and a neighborhood of $[(1, 0, 0)] \in X$, we get the assertion. $\square$

Since the S^1 -equivariant Euler class of an S^1 -module is given by the product of weights, we get the following.

Corollary 6.3. *The S^1 -equivariant Euler class of the tangent space at Z_λ is given by*

$$e(T_{Z_\lambda} X^{[n]}) = u^{2n} \prod_{s \in \lambda} (-\alpha a(s) - l(s) - 1) \cdot \prod_{s \in \lambda} (\alpha(a(s) + 1) + l(s)).$$

In particular, Z_λ is an isolated fixed point of S^1 -action and $e(T_{Z_\lambda} X^{[n]})$ is nonzero since $\alpha > 0$. In particular, we have

Corollary 6.4. *Let φ_λ be a nonzero element in $H_{S^1}^k(Z_\lambda)$. Then $\{i_{\lambda*} \varphi_\lambda \pmod{\text{Rad}\langle \cdot, \cdot \rangle}\}_\lambda$ forms an orthogonal basis of $H_{S^1, c}^{4n+k}(X^{[n]})/\text{Rad}\langle \cdot, \cdot \rangle$.*

Let us decompose the tangent space $T_{Z_\lambda} X^{[n]}$ as

$$N_{Z_\lambda}^{>0} \stackrel{\text{def.}}{=} \bigoplus_{p \in \mathbb{Z}} \bigoplus_{q > 0} H(p, q), \quad N_{Z_\lambda}^{\leq 0} \stackrel{\text{def.}}{=} \bigoplus_{p \in \mathbb{Z}} \bigoplus_{q \leq 0} H(p, q),$$

$$T_{Z_\lambda} X^{[n]} = N_{Z_\lambda}^{>0} \oplus N_{Z_\lambda}^{\leq 0}$$

Proposition 6.5. *The pull-back of the equivariant cohomology class $[L^\lambda C]_{S^1}$ by the inclusion $i_\lambda: Z_\lambda \rightarrow X^{[n]}$ is equal to the S^1 -equivariant Euler class of $N_{Z_\lambda}^{>0}$. It is given by*

$$i_\lambda^*[L^\lambda C]_{S^1} = e(N_{Z_\lambda}^{>0}) = u^n \prod_{s \in \lambda} (\alpha(a(s) + 1) + l(s)).$$

Proof. By the description (4.4) of $L^\lambda C$, it is nonsingular at Z_λ and its tangent space is equal to $N_{Z_\lambda}^{\leq 0}$. Hence the normal bundle is $N_{Z_\lambda}^{>0}$. Thus we have $i_\lambda^*[L^\lambda C]_{S^1} = e(N_{Z_\lambda}^{>0})$. The above formula follows from Lemma 6.2. $\square$

Theorem 6.6. *For each partition λ , let F_λ be the cohomology class*

$$(6.7) \quad i_{\lambda*} \left(\frac{u^n}{e(N_{Z_\lambda}^{\leq 0})} \right) = i_{\lambda*} \left(\frac{1}{\prod_{s \in \lambda} (-\alpha a(s) - l(s) - 1)} \right)$$

considered as an element in

$$H_{S^1, c}^{4n}(X^{[n]}) / \text{Rad} \langle \cdot, \cdot \rangle \cong H_{S^1}^{2n}(pt) \otimes H_c^{2n}(X^{[n]}) \xrightarrow[u^{-n} \otimes P.D.]{\cong} H_{2n}(X^{[n]}).$$

Then they satisfy

$$(6.8) \quad F_\lambda = [L^\lambda C] + \sum_{\mu < \lambda} u_{\lambda\mu}^{(\alpha)} [L^\mu C] \quad \text{for some } u_{\lambda\mu}^{(\alpha)},$$

$$(6.9) \quad \langle F_\lambda, F_\mu \rangle = 0 \quad \text{for } \lambda \neq \mu.$$

Corollary 6.10. *Under the identification $\bigoplus H_{2n}(X^{[n]}) \cong \Lambda \otimes \mathbb{C}$ given in Theorem 3.5, F_λ corresponds to the Jack's symmetric function $P_\lambda^{(\alpha)}$, where the parameter α is given by $-\langle C, C \rangle$ (see (1.1)).*

Proof of Theorem 6.6. The equation (6.9) follows from Theorem 6.1.

Let us take the generator $\varphi_\lambda \in H_{S^1}^0(Z_\lambda)$. By Corollary 6.4, we can write $[L^\lambda C]_{S^1}$ as a linear combination of $i_{\mu*} \varphi_\lambda$ modulo $\text{Rad} \langle \cdot, \cdot \rangle$. One can check that Z_μ is contained in $W^{\mu'}$. Hence $L^\lambda C = \text{Closure of } W^{\lambda'}$ (see Proposition 4.14) contains Z_μ only if $\mu \leq \lambda$ by (4.13). In particular, the pull-back $i_\mu^*[L^\lambda C]_{S^1} = 0$ unless $\mu \leq \lambda$. Hence we can write

$$[L^\lambda C]_{S^1} = \sum_{\mu \leq \lambda} v_{\lambda\mu}^{(\alpha)} i_{\mu*} \varphi_\lambda \quad \text{mod } \text{Rad} \langle \cdot, \cdot \rangle$$

for some $v_{\lambda\mu}^{(\alpha)}$.

In order to compute $v_{\lambda\lambda}^{(\alpha)}$, we consider the pull-back by i_λ^* . We have

$$\begin{aligned} i_\lambda^*[L^\lambda C]_{S^1} &= e(N_{\bar{Z}_\lambda}^{>0}) && \text{(by Proposition 6.5)} \\ &= \frac{e(T_{Z_\lambda} X^{[n]})}{e(N_{\bar{Z}_\lambda}^{\leq 0})} && \text{(since } T_{Z_\lambda} X^{[n]} = N_{\bar{Z}_\lambda}^{>0} \oplus N_{\bar{Z}_\lambda}^{\leq 0}) \\ &= \frac{1}{u^n} i_\lambda^* i_{\lambda*} \left(\frac{u^n}{e(N_{\bar{Z}_\lambda}^{\leq 0})} \right) && \text{(by the self-intersection formula).} \end{aligned}$$

Thus we get the assertion. $\square$

Finally note that the denominator in (6.7) coincides with the normalization factor (2.4) up to sign. Hence the integral form $J_\lambda^{(\alpha)}$ is an *integral* class. This shows that $J_\lambda^{(\alpha)}$ is expressed as a linear combination of m_λ with coefficients in $\mathbb{Z}$ for any positive integer α . Thus we have $J_\lambda^{(\alpha)} = \sum_{\mu < \lambda} \tilde{u}_{\lambda\mu}^{(\alpha)} m_\mu$ with $\tilde{u}_{\lambda\mu}^{(\alpha)} \in \mathbb{Z}[\alpha]$. This proves a part of Macdonald conjecture [17, 10.26?] mentioned in §2(i).

REFERENCES

- [1] M.F. Atiyah and R. Bott, *The moment map and equivariant cohomology*, Topology **23** (1984), 1–28.
- [2] H. Awata, Y. Matsuo, S. Odate, and J. Shiraishi, *Collective field theory, Calogero-Sutherland model and generalized matrix models*, Phys. Lett. **B347** (1995), 49–55.
- [3] R. Bott, *Vector fields and characteristic numbers*, Michigan Math. Jour. **14** (1967), 231–244.
- [4] J. Briançon, *Description de $\text{Hilb}^n \mathbb{C}\{x, y\}$* , Invent. Math. **41** (1977), 45–89.
- [5] N. Chriss and V. Ginzburg, “Representation theory and complex geometry (Geometric techniques in representation theory of reductive groups)”, Progress in Math. Birkhäuser, (to appear).
- [6] G. Ellingsrud and S.A. Strømme, *On the homology of the Hilbert scheme of points in the plane*, Invent. Math. **87** (1987), 343–352.
- [7] ———, *An intersection number for the punctual Hilbert scheme of a surface*, preprint, alg-geom/9603015.
- [8] J. Fogarty, *Algebraic families on an algebraic surface*, Amer. J. Math. **90** (1968), 511–521.
- [9] W. Fulton, “Intersection Theory”, A Series of Modern Surveys in Math. 2, Springer-Verlag, 1984.
- [10] L. Göttsche, *The Betti numbers of the Hilbert scheme of points on a smooth projective surface*, Math. Ann. **286** (1990), 193–207.
- [11] ———, *Hilbert schemes of zero-dimensional subschemes of smooth varieties*, Lecture Notes in Math. **1572**, Springer-Verlag, 1994.
- [12] L. Göttsche and W. Soergel, *Perverse sheaves and the cohomology of Hilbert schemes of smooth algebraic surfaces*, Math. Ann. **296** (1993), 235–245.
- [13] I. Grojnowski, *Instantons and affine algebras I: the Hilbert scheme and vertex operators*, Math. Res. Letters **3** (1996), 275–291.
- [14] A. Iarrobino, “Punctual Hilbert schemes”, Mem. Amer. Math. Soc. **188**, 1977.
- [15] F. Kirwan, “Cohomology of quotients in symplectic and algebraic geometry”, Mathematical Notes, Princeton Univ. Press, 1985.
- [16] F. Knop and S. Sahi, *A recursion and a combinatorial formula for Jack polynomials*, preprint, q-alg/9610016.
- [17] I.G. Macdonald, “Symmetric functions and Hall polynomials (2nd ed.)”, Oxford Math. Monographs, Oxford Univ. Press, 1995.

- [18] H. Nakajima, *Heisenberg algebra and Hilbert schemes of points on projective surfaces*, to appear in Ann. of Math.
- [19] ———, “Lectures on Hilbert schemes of points on surfaces”, to appear.
- [20] G. Wilson, *Collisions of Calogero-Moser particles and an adelic Grassmannian*, preprint.

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF TOKYO, KOMABA 3-8-1, MEGURO-KU,
TOKYO 153, JAPAN

E-mail address: nakajima@ms.u-tokyo.ac.jp